



ТАШКЕНТСКИЙ
ГОСУДАРСТВЕННЫЙ
СТОМАТОЛОГИЧЕСКИЙ
ИНСТИТУТ

MED UNION

МЕДИЦИНСКИЙ
НАУЧНО-ПРАКТИЧЕСКИЙ
ЖУРНАЛ

Volume 2, Issue 1

ISSN-2181-3183



ҚАДРЛИ ҲАМКАСБЛАР!

Маълумки, Ўзбекистонда ёшларга оид сиёсатга катта эътибор қаратилмоқда, айниқса, сўнгги йилларда Президентимиз ва ҳукуратимизнинг қатор меъёрий-ҳуқуқий ҳужжатлари қабул қилиниб, ёшларнинг илм-маърифат эгаллаши, меҳнат фаолияти ва бўш вақтини мазмунли ўтказиши учун кўпгина шарт-шароит яратишга хизмат қилмоқда.

Таклиф этилаётган «**MedUnion**» илмий-амалий журнали ёш олимлар, магистрлар, клиник ординаторлар, докторантлар, мустақил изланувчилар ва талабалар учун профессионал мулоқот майдони бўлиб хизмат қилади. Журнал электрон шаклда нашр этилади, чунки ушбу формат бир қатор афзалликларга эга: нашр этилган материаллар ҳажмига чекловлар олиб ташланади, муаллифдан ўқувчига бўлган йўл сезиларли даражада қисқаради, бу бизнинг динамик замонамизда жуда аҳамиятли, шунингдек ҳаражатлар ҳам анча камайтиради. Ҳар бир мақолага оригинал ДОИ рақами берилади.

Ушбу электрон илмий журналнинг мақсадлари:

- стоматология, умумий клиник, фундаментал фанлар, шунингдек, тиббиётда педагогика ва психология соҳасидаги замонавий тадқиқотларни ёритиш.
- ёш олимларнинг интеграциялашуви ва ушбу фанларнинг илмий ва амалиётчи мутахассислари ўртасидаги яқин ҳамкорлик.
- академик анъаналар давомийлигини сақлаш, илмий-педагогик кадрларни тарбиялаш.

Журналда ўзбек, рус ва инглиз тилларида ёш олимлар диссертацияларининг оригинал эмпирик тадқиқотлари ва умумий илмий-назарий мақолалар чоп этилади. Ишонаманки, ушбу журнал ҳақиқий мунозара майдонига айланади, илмий мулоқотни таъминлашга ёрдам беради, шунингдек, тиббиёт соҳасида янги илмий ва педагогик кадрларни тарбиялашга ўз хиссасини қўшади. Сизни ушбу лойиҳада турли материаллар муаллифи ва шарҳловчи сифатида иштирок этишга таклиф қиламиз.

Бош муҳаррир

Н. Ҳайдаров

Главный редактор:

Хайдаров Н.К. – д.м.н., ректор Ташкентского государственного стоматологического института (Узбекистан)

Заместитель главного редактора: Шомуродов К.Э.

– д.м.н., проректор по научной работе и инновациям ТГСИ (Узбекистан)

Ответственный секретарь:

Мун Т. О. – PhD, доцент ТГСИ (Узбекистан)

Редакционная коллегия:

Баймаков С.Р. – д.м.н., доцент ТГСИ (Узбекистан)
Амануллаев Р.А. – д.м.н., профессор ТГСИ (Узбекистан)
Ходжиметов А.А. – д.б.н., профессор ТГСИ (Узбекистан)
Мухамедов И.М. – д.м.н., профессор ТГСИ (Узбекистан)
Рустамова Х.Е. – д.м.н., профессор ТГСИ (Узбекистан)
Полатова Д.Ш. – д.м.н., профессор ТГСИ (Узбекистан)
Шамсиев Д.Ф. – д.м.н., профессор ТГСИ (Узбекистан)
Муртазаев С.С. – д.м.н., доцент ТГСИ (Узбекистан)
Вахидов У.Н. – д.м.н., доцент ТГСИ (Узбекистан)
Янгиева Н.Р. – д.м.н., доцент ТГСИ (Узбекистан)
Раимова М.М. – д.м.н., доцент ТГСИ (Узбекистан)
Салимов О.Р. – д.м.н., доцент ТГСИ (Узбекистан)
Хамдамов Б.З. – д.м.н., доцент БухГМИ (Узбекистан)
Собиров М.А. – д.м.н., доцент ТГСИ (Узбекистан)
Султонов Ш.Х. – д.м.н., доцент (Узбекистан)
Алимова Д.М. – д.м.н., доцент ТГСИ (Узбекистан)
Мирсалихова Ф.Л. – д.м.н., доцент ТГСИ (Узбекистан)
Пахомова Н.В. – к.м.н. доцент кафедры хирургической стоматологии и ЧЛХ ФГБОУ ВО ПСПбГМУ им. акад. И.П. Павлова (Россия)
Халматова М.А. – к.м.н., доцент ТГСИ (Узбекистан)
Махкамова Н.Э. – д.м.н., профессор ТГСИ (Узбекистан)
Нишанова А.А. – к.м.н., доцент ТГСИ (Узбекистан)
Туйчибаева Д.М. – д.м.н., доцент ТГСИ (Узбекистан)
Мухамедов Б.И. – к.м.н., доцент ТГСИ (Узбекистан)
Нугманова У.Т. – к.м.н., доцент ТГСИ (Узбекистан)
Дадабаева М.У. – к.м.н., доцент ТГСИ (Узбекистан)
Арипова Г.Э. – к.м.н., доцент ТГСИ (Узбекистан)
Шомухамедова Ф.А. – к.м.н., доцент ТГСИ (Узбекистан)
Махсумова С.С. – к.м.н., доцент ТГСИ (Узбекистан)
Содикова Х.К. – к.м.н., доцент ТГСИ (Узбекистан)
Алиева Н. М. – к.м.н., доцент ТГСИ (Узбекистан)
Скосырева О.В. – к.м.н., доцент ТГСИ (Узбекистан)

Chief editor:

Khaydarov N.K. – DSc, Rector of the Tashkent State Dental Institute (Uzbekistan)

Deputy Editor:

Shomurodov K.E. – DSc, Vice-Rector for Research and Innovation TSDI (Uzbekistan)

Executive assistant:

Mun T. O. – PhD, assoc.prof. of TSDI (Uzbekistan)

Editorial team:

Baymakov S.R. – DSc, assoc.prof. of TSDI (Uzbekistan)
Amanulaev R.A. – DSc, prof. of TSDI (Uzbekistan)
Khodjimetrov A.A. – DSc, prof. of TSDI (Uzbekistan)
Mukhamedov I.M. – DSc, prof. of TSDI (Uzbekistan)
Rustamova Kh.E. – DSc, prof. of TSDI (Uzbekistan)
Polatova D.Sh. – DSc, prof. of TSDI (Uzbekistan)
Shamsiev D.F. – DSc, prof. of TSDI (Uzbekistan)
Murtazaev S.S. – DSc, assoc. prof. of TSDI (Uzbekistan)
Vakhidov U.N. – DSc, assoc.prof. of TSDI (Uzbekistan)
Yangieva N.R. – DSc, assoc.prof. of TSDI (Uzbekistan)
Raimova M.M. – DSc, assoc.prof. of TSDI (Uzbekistan)
Salimov O.R. – DSc, assoc.prof. of TSDI (Uzbekistan)
Khamdamov B.Z. – DSc, assoc.prof. of BSMI (Uzbekistan)
Sobirov M.A. – DSc, assoc.prof. (Uzbekistan)
Sultanov Sh.Kh. – DSc, assoc.prof. of TSDI (Uzbekistan)
Alimova D.M. – DSc, assoc.prof. of TSDI (Uzbekistan)
Mirsalikhova F.L. – DSc, prof. of TSDI (Uzbekistan)
Pakhomova N. V. – PhD, assoc.prof. The First Saint Petersburg State medical university named after Academician Pavlov I.P. (Russia)
Khalmatova. M. A. – PhD, assoc.prof. of TSDI Uzbekistan)
Makhkamova N.E. – DSc, prof. of TSDI Uzbekistan)
Nishanova A.A. – PhD, assoc.prof. of TSDI (Uzbekistan)
Tuychibaeva D.M. – DSc, assoc.prof. of TSDI (Uzbekistan)
Mukhamedov B.I. – PhD, assoc.prof. of TSDI (Uzbekistan)
Nugmanova U.T. – PhD, assoc.prof. of TSDI (Uzbekistan)
Dadabaeva M.U. – PhD, assoc.prof. of TSDI (Uzbekistan)
Aripova G.E. – PhD, assoc.prof. of TSDI (Uzbekistan)
Shomukhamedova F.A. – PhD, assoc.prof. of TSDI (Uzbekistan)
Makhsumov S.S. – PhD, assoc.prof. of TSDI (Uzbekistan)
Sodikova Kh.K. – PhD, assoc.prof. of TSDI (Uzbekistan)
Alieva N.M. – PhD, assoc.prof. of TSDI (Uzbekistan)

Реймназарова Г.Д. – к.м.н., доцент ТГСИ (Узбекистан)
Аляви С. Ф. – к.м.н., доцент ТГСИ (Узбекистан)
Каримова М.У. - к.м.н., доцент ТГСИ (Узбекистан)
Акрамова Л.Ю. – к.п.н., доцент ТГСИ (Узбекистан)
Бабакулов Ш. Х.- PhD, доцент ТГСИ (Узбекистан)
Хасанова Л.Э.- д.м.н., профессор ТГСИ (Узбекистан)
Балтабаев У.А.- д.х.н., профессор ТГСИ (Узбекистан)
Буранова Д.Д.-к.м.н., доцент ТГСИ (Узбекистан)
Исраилова М.Н.- PhD, доцент ТГСИ (Узбекистан)
Хикметов Б.А. - PhD, доцент ТГСИ (Узбекистан)
Абдукодиров Э.И. - PhD, доцент ТГСИ (Узбекистан)

Редакционный совет:

Jaе Hoon Lee – д.м.н., профессор Университет Ёнсей (Южная Корея)
Kavinda Sudharaka Tissera – PhD, Университет Рухуна (Шри Ланка)
Ермак О.А. – к.м.н., доцент Белорусской медицинской академии последиplomного образования (Беларусь)
Бекжанова О.Е. – д.м.н., профессор ТГСИ (Узбекистан)
Даминова Л.Т.– д.м.н., профессор ТГСИ (Узбекистан)
Дусмухамедов М.З. – д.м.н., профессор ТГСИ (Узбекистан)
Азизов Б.С. – д.м.н., доцент ТГСИ (Узбекистан)
Пулатова Б.Д.– д.м.н., доцент ТГСИ (Узбекистан)

Дизайн и технические работы:

Мирхайидов М.М.
Жураев Б.Н.
Мусаев Ш.Ш.

Skosireva O.V. – PhD, assoc.prof. of TSDI (Uzbekistan)
Reimnazarov G.D. – PhD, assoc.prof. of TSDI (Uzbekistan)
Alyavi S.F. – PhD., assoc.prof. of TSDI (Uzbekistan)
Karimova M.U. - PhD, assoc.prof. of TSDI (Uzbekistan)
Akramova L.Yu. – PhD, assoc.prof. of TSDI (Uzbekistan)
Babakulov Sh.Kh. - PhD, assoc.prof. of TSDI (Uzbekistan)
Khasanova L.E.- DSc, assoc.prof. of TSDI (Uzbekistan)
Baltabayev U.A. - DSc, professor in TSDI (Uzbekistan)
Buranova D. D.- PhD, assoc.prof. of TSDI (Uzbekistan)
Israilova M. N.- PhD, assoc.prof. of TSDI (Uzbekistan)
Khikmetov B.A.- PhD, assoc.prof. of TSDI (Uzbekistan)
Abdukodirov E.I.- PhD, assoc.prof. of TSDI (Uzbekistan)

Editorial Council:

Jaе Hoon Lee – DSc, Prof. of Yonsei University (South Korea)
Kavinda Sudharaka Tissera – PhD, Ruhuna University (Sri Lanka)
Ermak O.A. – PhD, assoc.prof. of the Belarusian Medical Academy of Postgraduate Education (Belarus)
Bekjanova O.E. – DSc, prof. in TSDI (Uzbekistan)
Daminova L.T. – DSc, prof.in TSDI (Uzbekistan)
Dusmukhamedov M. Z. – DSc, prof. in TSDI (Uzbekistan)
Azizov B. S. – DSc, assoc.prof. of TSDI (Uzbekistan)
Pulatov B. D.– DSc, assoc.prof. of TSDI (Uzbekistan)

Design and technical work:

Mirkhayidov M.M.
Juraev B.N.
Musaev Sh.Sh.

Содержание

1. Абдуқодиров Э.И., Абдуллаева М.Б., Чориева Ф.Н., Дустмуродов О.Ҳ., Ходжаева К. А. Тригеминал невралгияларда замонавий даво усулларини қўллаш	6-12
2. Абдурахмонов И.Р., Шамсиев Дж.Ф. Бош мия фалажи фондаги параназал синуситларни даволашда ўзига хос ёндашиш	13-25
3. Абдусаматова И. И., Шамсиев Д.Ф., Тастанова Г.Э. Буруннинг нафас ўтказувчанлиги бузилишида механорецепторларнинг клиник аҳамияти (адабиёт шарҳи)	26-34
4. Азизов Б.С., Нурматова И.Б., Саиткулов Э.Х. Клинический случай туберозного склероза	35-44
5. Акрамова Л.Ю., Акрамов Н.М. О гуманитарной составляющей в современной подготовке врача	45-51
6. Арифов С.С., Тухтаев М.Б. Актуальные вопросы диагностики болезни Меньера	52-61
7. Ахмадалиев Н.Н., Режаббоева Н.Л. Вирусли гепатитларда холестатик синдром ва апоптознинг биокимёвий маркерлари	62-68
8. Babayev J., Kurbanniyazova Sh., Sultonov Sh. O'smirlar ruhiy salomatligi: raqamli dunyoda xavf va imkoniyatlar	69-73
9. Билял Н.М., Арипова Г.Э. Особенности лечения дистального прикуса аппаратом твин-блок	74-80
10. Газарян Л.Г., Ордиянц И.М., Савичева А.М., Мухаммаджанова М.О. Аминокислоты и их метаболиты как маркеры прогнозирования акушерских и перинатальных осложнений при гестационном сахарном диабете	81-87
11. Ибрагимов Д.Д., Мардонова Н.П., Исматов Н.С., Кучкоров Ф.Ш., Жағ кисталарини даволашда тромбоцитлар билан тўйинган фибриннинг қўллаш авзаллиги	88-93
12. Ибрагимов Д.Д., Отабоев Р.Ш. Профилактика деформации альвеолярного отростка челюстей с применением костного регенератора стимул	94-99
13. Каюмова С.А., Туйчибаева Д.М., Урманова Ф.М. Анализ эффективности применения ксенотрансплантата в хирургическом лечении пациентов с возрастной макулярной дегенерацией	100-105
14. Ким А.А., Туйчибаева Д.М. Распространенность и факторы риска кератоконуса (обзор литературы)	106-114
15. Khvan O.I., Don A.N. Spleen injuries during blunt trauma of the abdominal	115-119
16. Мамырбекова С.А., Раушанова А.М., Рустамова Х.Е., Нурмаматова К.Ч. Казахский национальный университет им. аль-фараби – путь к элитному образованию и межеуззовским сотрудничествам	120-124
17. Мизомов Л.С., Азимов А.М. Тепловизионное исследование лица взрослых больных острым одонтогенным остеомиелитом челюстей	125-131
18. Мирсалихова Ф.Л., Эронов Ё.Қ. Имконияти чекланган болаларда пародонт касалликларни ташхислаш ва стоматологик текширув усуллари	132-138

19. Мухаммаджанова М., Курбанов А.К., Буранова Д.Д., Хасанова Х.Ж., Акбарова Г.П. Коррекция артериальной гипертензии у пациентов пожилого возраста	139-154
20. Мухутдинов Э.Р. Атабекова Ш.Н., Изменения стоматологического статуса у спортсменов при синдроме перетренированности организма	155-160
21. Nurmatov U.B., Nurmatova N.F., Baxodirova I.U. Study of the microflora and ph of the skin environment in patients with zoonanthropontic trichophytosis of the pubic region	161-169
22. Орипова А.А., Шарипова А.У. Состояние верхнечелюстной пазухи у пациентов с «перелом скуловой кости»	170-176
23. Otamuratov R.U. Internet ijtimoiy tarmoqlari foydalanuvchilari faoliyatining o'ziga xos xususiyatlari bilan shaxsiy xususiyatlarning aloqasi	177-182
24. Раимова М.М., Мухамедсаидова И.А., Бабашева Д.Р. Безовта оёқлар синдроми клиник кўринишлари, диагностика ва давоси	183-190
25. Рахимова Г.Ш. Шамсиев Д.Ф. Современные принципы профилактики и лечения беременных с аллергическим ринитом	191-201
26. Рустамова Х.Е., Нурмаматова К.Ч., Рахимжонов А.А., Рустамова М.А. Охрана здоровья женщин как медико-социальная проблема	202-207
27. Сафарова Э.М., Юнусов Р.А., Рахимова И.И. Суицид как медико-социальный аспект: сравнительный анализ суицида	208-215
28. Тахирова К.А., Кадирова М.Н., Разакова Н.Б. Проблема сочетанной патологии полости рта и органов пищеварения у подростков	216-221
29. Туйчибаева Д.М., Ким А.А. Современные подходы в лечении кератоконуса в разных странах мира (обзор литературы)	222-240
30. Тилляшайхов М.Н, Бойко Е.В., Саламов М.С., Юсупов Ш.Х., Бобокулов Ш.Х. Метастатический рак простаты - рандомизированное исследование ципротерон ацетата в сравнении с флутамидом	241-246
31. Усманбекова Г.К. Системный анализ уровня подготовки последипломного образования среднего медицинского персонала стоматологических учреждений	247-253
32. Усманова Н.А., Махкамова Н.Э. Обзор анатомических и морфологических данных о слуховой трубе, возможных причинах ее дисфункций	254-260
33. Xusanходjaeva F.T., Daminova K.M. Lyupus nefrit rivojlanishning patogenetik mexanizmlari	261-270
34. Xusanходjaeva F.T., Xasanova X.Dj., Buranova D.Dj., Salyamova F.E. Zamonaviy gadgetlarning talabalarning o'quv jarayoniga salbiy ta'siri	271-276
35. Шукпаров А.Б., Шомуродов К.Э. Результаты направленной костной регенерации после предварительного расширения мягких тканей	277-285
36. Якубходжаева М.Р. Синтез и биологическая активность 1,2,3-триазолов на основе ацетеленсодержащих производных карбаматов	286-291

УДК: 616.992:616.992.28

STUDY OF THE MICROFLORA AND PH OF THE SKIN ENVIRONMENT IN PATIENTS WITH ZOOANTHROPONOTIC TRICHOPHYTOSIS OF THE PUBIC REGION

¹Nurmatov U.B., ²Nurmatova N. F., ³Baxodirova I.U.

¹Tashkent State Dental Institute

²Tashkent medical academy,

³Tashkent pediatric medical institute
Tashkent, Uzbekistan

SUMMARY

The article describes the Study of the microflora and pH of the skin environment in patients with zooanthroponotic trichophytosis.

Key words: Zooanthroponotic trichophytosis and features of microbiology.

Аннотация

В статье описано исследование микрофлоры и pH кожной среды у больных зооантропонозной трихофитией.

Ключевые слова: зооантропонозная трихофития и особенности микробиологии.

Annotatsiya

Maqolada zooantroponotik trixofitozli bemorlarda mikroflora va teri muhitining pH ni o'rganish tasvirlangan.

Kalit so'zlar: zooantroponotik trixofitoz va mikrobiologiyaning xususiyatlari.

The relevance of the work. Serious ecological changes taking place on the planet in recent decades could not but affect the state of microecology and immunoreactivity of modern man. This is one of the global reasons for the increase in the number of fungal diseases. In addition, outbreaks of zooanthroponotic dermatomycosis are also caused by a change in the properties of the pathogens themselves - the emergence of new strains of fungi, an increase in their pathogenicity and contagiousness, a violation of the structure and boundaries of nosoareals. It should be noted that today in Uzbekistan, favus has been eliminated almost everywhere, and the incidence of anthroponotic dermatomycosis has been sharply reduced. At the same time,

zooanthroponotic trichophytosis has a wide and widespread distribution throughout the territory of the Republic.

Zooanthroponotic trichophytosis is a natural focal infection, the epidemic features and the frequency of outbreaks of which depend on geographical, environmental and a number of other factors [1]. In modern conditions, trichophytosis in the Republic of Uzbekistan has a number of characteristic features: the species composition of pathogens has changed (the main pathogen is trichophyton faviform, whose share in the total structure of trichophytosis pathogens is 80-85%). The age composition of patients also changed (at the end of the 90s, 86% of patients with trichophytosis were children of preschool and primary school age, and by 2003, 61%

of patients were over 15 years old), atypical localizations of trichophytosis lesions appeared, including the pubic area, genitals, buttocks [1,3].

Since 2002, registration of patients with zoonanthroponotic trichophytosis with localization of the process in the genital area has begun. In all regions of the Republic, cases of registration of zoonanthroponotic trichophytosis among the adult population with a predominant lesion of the pubic and inguinal region have become more frequent, and the vast majority of patients indicate the sexual route of infection. There is a direct relationship between the frequency of infection with sexually transmitted infections (STIs) and the occurrence of pubic trichophytosis. Given the steady increase in the number of STIs and the complications they cause, we should expect an outbreak of morbidity and pubic trichophytosis.

In this connection, knowledge of the distribution features, conditions of clinical manifestation and characteristics of the pathogen, the development of adequate methods of treatment will allow influencing the main links of the epidemic process.

The degree of knowledge of the problem. Despite ongoing measures to prevent fungal diseases of the skin around the world, there is an increase in infection with this infection. Thus, in the Republic of Kazakhstan, the incidence of fungal infections of the skin in recent years has increased from 3756 cases in 1999 to 5401 cases in 2003 with a predominant lesion of the urban (2412 people) and male (3310 people) population [1]. The study of the incidence of zoonanthroponotic trichophytosis in different regions of the CIS made it possible to find out that against the background of a significant decrease in the total number of patients with fungal

diseases, it remains at a fairly high level, amounting to 95% in rural areas. [1,2]

In the Republic of Uzbekistan, there is also a widespread increase in the incidence of zoonanthroponotic trichophytosis. Since a dry, hot climate prevails on the territory of Uzbekistan, the highest incidence of trichophytosis occurs in the summer period of the year. This is due to the fact that skin moist from excess sweat, often macerated stratum corneum of the epidermis, a shift in the pH of the skin and sweat towards an alkaline reaction contributes to the penetration of pathogens into the skin and the development of fungal infections in it [1,5].

Currently, zoonanthroponotic trichophytosis is characterized by a pronounced clinical diversity. The pathogenesis of the development of clinical forms of mycosis can be influenced by the characteristics of the nature and intensity of the body's immune response to the introduction of the pathogen, the degree of their virulence and pathogenicity, as well as disorders of the immunogenesis system. Numerous clinical studies have shown that at the present stage of development of mycology, a change in the spectrum of mycosis pathogens, the development of fungal-bacterial associations entail not only a change in the characteristic clinical manifestations of diseases, but also, importantly, the development of resistance to the antifungal agents used [1,2]. However, a comprehensive study of the most significant factors influencing the development of pubic trichophytosis, the state of the body's immune reactivity, the study of the nature of the accompanying microflora and skin pH in combination with various clinical forms of trichophytosis in Uzbekistan was not carried out.

It is known that against the background of an imbalance in the immune system and a

decrease in nonspecific protective factors of the body, the microbial landscape of the body as a whole, and in particular on the skin, changes. In the foci of zooanthroponotic trichophytosis, often associated microflora complicates the mycotic process, which changes the clinical course of dermatosis. Moreover, the role of secondary microflora in the foci of trichophytosis and the mechanism of its influence on the clinic of the disease have not been studied enough. In addition, the pH environment of the skin plays an important role in the development of this pathology, which contributes to the creation of favorable or unfavorable conditions for the development of various microorganisms and the development of their dysbiosis.

Purpose of the study. To study the state of the accompanying microflora and the pH of the skin environment in the lesions in patients with pubic trichophytosis.

Material and methods. The state of the microflora and pH of the skin environment with zooanthroponotic trichophytosis of the pubic area was studied by us in 84 patients in comparison with the data of 20 practically healthy individuals (control group). Among the examined patients with zooanthroponotic trichophytosis of the pubic region, 20 patients had a superficial-spotted form, 22 had an infiltrative form, and 42 had an infiltrative-suppurative form of the disease.

In the study of skin washings from the pubic area in the control group, the content of microbes ranged from 2.0 to 5.0 CFU/cm². In this group, microbiological detection of various microbes was observed in 4 out of 20 individuals. As can be seen from Table 1, the microbial landscape of the skin of the pubic region in apparently healthy individuals includes *Staphylococcus epidermidis*, *aureus*, and *Candida*.

Table 1.

The state of the microflora of the pubic area in the control group (n=20).

Type of microflora	The number of microbes per 1 cm ² of skin (CFU / cm ²)
<i>Staphylococcus epidermidis</i>	5,0 ± 0,3
<i>Staphylococcus aureus</i>	2,0 ± 0,2
<i>Streptococcus haemolyticus</i>	-
<i>Clostridium perfringens</i>	-
<i>Candida</i>	3,0 ± 0,5
<i>Klebsiella</i>	-
<i>Enterococcus</i>	-

The study of the state of the microflora of the skin of the pubic region in 84 patients with zooanthroponotic trichophytosis showed that the cultural study of skin washings revealed different microflora from the lesion. Among the identified microorganisms from the lesions of

trichophytosis, *Staphylococcus epidermidis* was sown in 36 (42.9%), *Staphylococcus aureus* - in 27 (32.1%), *Clostridium perfringens* - in 11 (13.1%), *Candida* - in 6 (7, 1%), *Klebsiella* - in 2 (2.4%) cases, and *Enterococcus* and *Streptococcus haemolyticus* - in 1 (2.4%) case.

Table 2.

The state of the microflora of the pubic area in the control group (n=84).

Type of microflora	Total number of microbes per 1 cm ² of skin (CFU/cm ²)
Staphylococcus epidermidis	109,0 ± 5,2
Staphylococcus aureus	97,0 ± 4,3
Streptococcus haemolyticus	14,0 ± 1,2
Clostridium perfringens	22,3 ± 2,3
Candida	4,0 ± 0,2
Klebsiella	10,7 ± 4,2
Enterococcus	8,1 ± 0,4

Along with this, when examining the microflora in the lesions of patients with trichophytosis of the pubic region, changes were observed both in qualitative and quantitative characteristics in relation to the control group (Table 2). As can be seen from the table, in the focus of trichophytosis of the pubic region, there is a qualitative predominance of microflora (Streptococcus haemolyticus, Clostridium perfringens, Klebsiella, Enterococcus), which is not a normoflora in the microbial landscape of the skin of healthy individuals. In a place with this, their number increases, for example; St. epidermidis in the control group was 5.0 ± 0.3 CFU/cm², and in the focus of trichophytosis of the pubic region, this figure increased to 109.0 ± 5.2 CFU/cm² ($p < 0.001$), an analogous picture was observed with St. aureus 2.0 ± 0.2 and 97.0 ± 4.3 CFU/cm², respectively ($p < 0.001$). In the quantitative indicator of fungi of the genus Candida, there was a slight upward trend (3.0 ± 0.5 and 4.0 ± 0.2 CFU / cm², respectively), perhaps small quantitative fluctuations are associated with the antagonistic properties of fungi of the genus

Candida and Trichophyton, manifested in growth suppression fungi of the genus Candida last [3].

Considering the revealed imbalance in the microbiological landscape of the skin of the pubic area in case of trichophyton damage, it seemed very interesting to study the state of the microflora in the lesions depending on the clinical forms of zooanthroponotic trichophytosis of the pubic area. Studies conducted in 20 patients with a superficially spotty form of zooanthroponous trichophytosis of the pubic region showed that when sowing skin washings of Staphylococcus epidermidis was sown in 9 (45.0%) cases, Staphylococcus aureus - in 4 (20.0%), Clostridium perfringens - in 2 (10.0%) and Candida - in 5 (25.0%) cases.

The study of the microbiological landscape in case of damage to the surface-spotted form of zooanthroponous trichophytosis of the pubic region showed a quantitative increase in St. epidermidis compared with the control group 5.0 ± 0.3 CFU/cm² versus 19.0 ± 4.8 CFU/cm², St. aureus 2.0 ± 0.2 CFU/cm² and 14.0 ± 6.4 CFU/cm², respectively (Table 3).

Table 3.

The state of the microflora in the focus of zooanthroponotic trichophytosis of the pubic region depending on the clinical forms of the disease

Type of microflora	The number of microbes per 1 cm ² of skin (CFU / cm ²)		
	Superficial spotted form (n=20)	Infiltrative form (n=22)	Infiltrative suppurative form (n=42)
Staphylococcus epidermidis	19,0 ± 4,8	35,4± 7,1	54,6 ± 3,5
Staphylococcus aureus	14,0 ± 6,4	32,0 ± 9,2	51,0 ± 6,0
Streptococcus haemolyticus	-	-	14,0
Clostridium perfringens	4,3 ± 3,0	6,4 ± 4,4	11,6 ± 3,7
Candida	2,0 ± 0,8	2,0	-
Klebsiella	-	-	6,0
Enterococcus	-	-	8,1 ± 5,7

The study of microflora inoculation in patients with infiltrative form of zooanthroponous trichophytosis showed that among patients with this clinical form of the disease, Staphylococcus epidermidis was found in 11 (50.0%), Staphylococcus aureus - in 8 (36.4%), Clostridium perfringens - in 2 (9.1%) and Candida - in 1 (4.5%) cases.

The study of the microbiological landscape with the defeat of the infiltrative form of zooanthroponous trichophytosis of the pubic region revealed both a qualitative and quantitative increase in various types of microorganisms. In quantitative terms, St. epidermidis in the infiltrative form of the disease was 35.4± 7.1 CFU/cm², which prevailed in comparison with the control group 5.0 ± 0.3 CFU / cm², and in relation (19.0 ± 4.8 CFU / cm²) to indicators for the surface-spotted form of zooanthroponotic trichophytosis. A similar pattern of quantitative increase was observed for Staphylococcus aureus, which amounted to 32.0 ± 9.2 CFU/cm². On the contrary, in the

infiltrative form of the disease, a decrease in the inoculation of fungi of the genus Candida was noted, which was detected in this group only in one case. It is especially important to note that with a change in the form of zooanthroponotic trichophytosis of the pubic region, there is a qualitative change in the composition of microbes in the lesion, which is expressed by the detection of such a microorganism as Clostridium perfringens, which was identified in our studies. As is known, these microorganisms are extremely sensitive to drying and do not multiply on healthy skin, however, according to the scientific literature, a person can be its carrier on the skin in 40% of cases [4]. This microorganism is able to infect wound surfaces and cause a destructive abscess, where the products of its metabolism can cause toxemia, which is probably what characterizes the violations of the general condition of patients (weakness, fever, etc.) with deep infiltrative-suppurative forms of the disease.

Next, we studied the microbiological state of the skin of the pubic region with a deep (infiltrative-suppurative) form of zooanthroponotic trichophytosis. The results of cultural studies showed an increase in the qualitative indicator with the seeding of lesions with new types of microbes such as *Klebsiella* in 2 (4.8%) cases, as well as *Enterococcus* and *Streptococcus haemolyticus* in 1 (4.8%) case (Fig. 3.9).

Along with them, the quantitative ratio of *Staphylococcus epidermidis* increased - 54.6 ± 3.5 CFU / cm², *Staphylococcus aureus* - 51.0 ± 6.0 CFU / cm², *Clostridium perfringens* - 11.6 ± 3.7 CFU / cm² in the lesion, compared with the superficial - spotted and infiltrative forms of the disease. The study of microflora inoculation in patients with infiltrative-suppurative form of zooanthroponous trichophytosis of the pubic region showed that among patients *Staphylococcus epidermidis* was found in 16 (38.0%), *Staphylococcus aureus* - in 15 (35.7%), *Clostridium perfringens* - in 7 (16, 7%), *Klebsiella* - in 2 (4.8%) cases, and *Enterococcus* and *Streptococcus haemolyticus* in 1 (2.4%) case (Fig. 4). And fungi of the genus *Candida* were not detected in this group of patients, which is consistent with the latest literature data on antagonistic properties with fungi of the genus *Trichophyton*, which, in quantitative terms, becomes larger in the lesions of the infiltrative and infiltrative-suppurative form of zooanthroponic trichophytosis of the pubic region.

The results of the study showed that in all clinical forms of pubic trichophytosis, there is a combination of concomitant microflora with the causative agent of the underlying disease, which is characterized by both a qualitative and quantitative increase in it in the lesion in direct proportion to the clinical forms of the disease.

It is especially important to note that with the transition to more severe forms of the disease, there is a large contamination of the lesions with various types of microorganisms, especially the detection of anaerobic species such as *Clostridium*, *Klebsiella*, which are unsuitable for reproduction on the surface of the skin. However, given the fact that the dermis actively metabolizes oxygen diffusing through the epidermis and taking into account the characteristics of the pubic and perigenital zones, such as relative humidity (over 90%) and the effect of occlusion, which leads to an increase in the pH of the environment in these areas of the skin, various microorganisms (aerobes and anaerobes) find favorable conditions for survival and generation. Identical conditions (humidity 97%) are favorable for growth and reproduction of trichophyton dermatomycetes [4,5]. Optimal conditions, the symbiosis of these microorganisms, violations of the functions of the immune system, which cannot adequately control and fight secondary infections, are probably the main factors contributing to the development of atypical localization of trichophytosis and aggravation of the course of mycotic dermatosis with a deeper lesion of the skin and the development of an infiltrative-purulent process with trichophytosis of the pubic region.

In this connection, in further studies, we studied the pH level of the skin environment in patients with zooanthroponotic trichophytosis of the pubic region.

A comparative study of the pH of the environment of the affected area of the skin of the pubic area in patients with trichophytosis was carried out in 84 patients, depending on the clinical forms of the disease, in comparison with the data of 20 practically healthy individuals (control group). As is known from the scientific

literature, the pH environment of the pubic region, perineum and perigenital zones has a neutral pH environment, while in other parts of the skin, these values average 5.4 [4,5].

The results of our studies of the pH of the medium in the control group are consistent with the scientific literature and average

6.98 ± 0.10 . Studies of the pH of the environment in patients with zooanthroponotic trichophytosis of the pubic area in the lesion, significantly increase compared with the data of the control group ($p < 0.001$) and average 7.52 ± 0.07 with 6.98 ± 0.10 in the control (table 4).

Table 4.
pH index of the skin environment in patients with various clinical forms of trichophytosis of the pubic area ($M \pm m$)

Clinical forms	The number of examined persons	pH index of the skin environment
Control	20	$6,98 \pm 0,10$
In the lesion focus of patients with trichophytosis	84	$7,52 \pm 0,07^*$
Superficial spotty	20	$7,50 \pm 0,11^*$
Infiltrative	22	$7,56 \pm 0,09^*$
Infiltrative suppurative	42	$7,50 \pm 0,13^*$

Note: p is the reliability of the data in relation to the control. * - $p < 0.001$

Analysis of changes in the pH value of the skin environment in patients with zooanthroponotic trichophytosis, depending on the clinical forms, revealed that in all the examined groups of patients with various clinical forms of the disease, the same statistically significant increase in the pH of the skin environment in the lesions was observed in relation to the indicators of healthy individuals ($p < 0.001$). This indicates that in this pathology there is a violation of the pH value of the skin environment with its shift to the alkaline side. The indicators themselves did not differ significantly in different clinical forms of the disease.

Based on the research results and scientific literature data, it should be noted that the

increase in the number of cases of zooanthroponotic trichophytosis of the pubic area, along with the above factors, is characterized by an increase in the pH values of the skin environment of the pubic area to the alkaline side, which also increases by 1.5 units in hyperhidrosis, which is often created by a greenhouse zone effect. [5,6]. In addition, the alkaline environment, being favorable for the generation of secondary microflora, promotes loosening of the epidermis and facilitates a deeper penetration of trichophyton and secondary microflora into the skin, creating favorable soil for the development of deep forms (infiltrative and infiltrative-suppurative) of the disease with the development of an infiltrative-purulent

process. with zooanthroponotic trichophytosis of the pubic region.

Thus, the studies showed an abundance of microbial landscape in the foci of zooanthroponotic trichophytosis of the pubic region with the detection of *Staphylococcus epidermidis* - in 42.9%, *Staphylococcus aureus* - 32.1%, *Clostridium perfringens* - 13.1%, *Candida* - in 7.1%, *Klebsiella*- 2.4%, *Enterococcus* and *Streptococcus haemolyticus* - in 2.4% of cases. Which were characterized by both qualitative and quantitative increase in the

lesion in direct proportion to the clinical forms of the disease. In patients with zooanthroponotic trichophytosis of the pubic region, the disease occurs against the background of a violation of the pH value of the skin environment with a shift to the alkaline side, which contributes to the creation of favorable conditions for the reproduction of various microorganisms with the development of their dysbiosis and complicates the clinical course of the mycotic process.

Литература / References

1. Арифов С.С., Абидова З.М., Лукьянова А.С. Зооантропонозная трихофития атипичной локализации среди взрослого контингента //Клин. дерматол. и венерол. –2003. –№3. –С.23-26.
2. Акимов В.А. Изменение показателей прямого счета бактерий в культурах и прудовой воде при различном pH //Микробиология. –1975. –Т. 44. –Вып.1. – С.163-165.
3. Мусаева Н.Ш. Оценка диагностических характеристик кожи при зооантропонозной трихофитии и совершенствование ее терапии: Автореф. Дис. ... канд. мед. наук. – Т., 2006.- 17с.),
4. Нобл У.К. Микробиология человека. – М: «Медицина», 1986. – 493 с.).
5. Т.Г. Терешкова Г.М., Калачева Г.С., Сальников М.В. Влияние pH среды на рост и физиологию водородокисляющих бактерий /Волова //Микробиология. –1986. –Т.55. –Вып.5. – С.745-749.
6. Behrendt H., Green M. Patterns of Skin pH from Birth Through Adolescence. Springfield, Illinois: Thomas, 1971.
7. ИБ Нурматова, БС Азизов «Фунготербин» в лечении микозов кожи, Фергана, «Перспективы развития новых технологий в диагностике дерматовенерологии и дерматоонкопатологии» 2022, май, сборник статей и тезисов конференции, Стр.106
<http://repository.tma.uz/xmlui/handle/1/3249>
8. Азизов Б., Исмаилова Г., Набиев Ф., Нурматова И., Усманов А. Клиническая характеристика больных ВИЧ/СПИДом // Новости дерматовенерологии и репродуктивного здоровья, 2013 год, №3, Стр. 46-49
<https://scholar.google.ru/scholar?oi=bibs&cluster=18328685665296077428&btnI=1&hl=ru>
9. MM Abdurakhmanov, NS Akramova, IB Nurmatova, BS Azizov// Роль хеликобактерной инфекции в развитии функциональной диспепсии и вульгарных угрей// Likars' ka sprava, 45–53-45–53.
https://scholar.google.ru/citations?view_op=view_citation&hl=ru&user=j5w7q9AAAAAJ&citation_for_view=j5w7q9AAAAAJ:KxtntwgDAa4C

10. ФИ Иноятова, ГЗ Нурматова Н.Ф, Иногамова// Роль лямблиозной инвазии в течении хронической вирусной патологии у детей. Совершенствование диагностики и лечения
https://scholar.google.ru/citations?view_op=view_citation&hl=ru&user=NSekut8AAAA

J&citation_for_view=NSekut8AAAAJ:D_slNldO8mEC

11. ФИ Иноятова, ШУ Абдумаджидова, НФ Нурматова // Дифференциально-диагностические информативные критерии хронического вирусного гепатита D у детей